APS/123-QED

arXiv:0712.0547v3 [astro-ph] 11 Jan 2008

Far Fields and Dark Matter

Andrea Carati[*]
Department of Mathematics, University of Milano
Via Saldini 50, 20133 Milano, Italy

Sergio L. Cacciatori[†]
Department of Physical and Mathematical Sciences, Insubria University
Via Valleggio 11, I-22100 Como, Italy

Luigi Galgani[‡]
Department of Mathematics, University of Milano
Via Saldini 50, 20133 Milano, Italy

(Dated: February 2, 2008)

We give an estimate of the gravitational field of force exerted on a test particle by the far galaxies, in the frame of the weak field approximation. In virtue of Hubble's law, the action of the far matter turns out to be non negligible, and even the dominant one. A nonvanishing contribution is obtained only if the discrete and fractal nature of the matter distribution is taken into account. The force per unit mass acting on a test particle is found to be of the order of $0.2\, cH_0$, where c is the speed of light and H_0 the present value of Hubble's constant.

PACS numbers: 95.35.+d, 98.80.-k

The main idea underlying the present paper comes from a critical reading of the first work in which the existence of dark matter was proposed, namely the work of Zwicky [1] in which the virial theorem was applied to the Coma cluster. The point concerns the role of the forces due to the external matter. Let us recall that in the first use of the virial theorem, namely that of Clausius for a gas enclosed in a vessel, the main contribution to the virial was coming from the external forces (the pressure of the walls), because the contribution of the internal forces was either vanishing (for a perfect gas) or negligible. In the case of Coma, the contribution of the internal forces was still found by Zwicky to be negligible, but the contribution of the external galaxies was neither taken into account, nor even mentioned at all, perhaps because one could not conceive how it may act as a pressure. The missing contribution to the virial was thus attributed to some hypothetical interior dark matter. In the present letter we show how it is conceivable that the external matter (actually, the far one) may produce a pressure, and we also give a rough estimate of its contribution to the virial, which appears to be in rather good agreement with the observations.

The first point we make is that, as the forces on a test particle depend not only on the positions of the galaxies, but also on their velocities and accelerations, due to Hubble's law the dominating contribution comes from the far matter. This comes about as follows. From the point of view of general relativity, in the weak–field approximation the problem of estimating the force (per unit mass) on a test particle amounts to writing down the equations for the geodesic motion when the metric tensor $g_{\mu\nu}$ is a solution of the Einstein equation with the external galaxies as sources. Writing the metric tensor as a perturbation of the Lorentzian background $\eta_{\mu\nu}$, namely, as $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, the perturbation $h_{\mu\nu}$ turns out to be a solution of the wave equation, so that its components are analogous to the familiar retarded potentials of electrodynamics (although relevant differences exist between the two cases, as particularly emphasized by Zeldovich and Novikov [2]). In fact one finds

$$h_{\mu\nu} = \frac{-2\pi G}{c^4} \sum_j M_j \frac{1}{\gamma_j} \left. \frac{2\dot{q}_\mu^{(j)}\dot{q}_\nu^{(j)} - c^2\eta_{\mu\nu}}{|\mathbf{x}-\mathbf{q}^{(j)}|}\right|_{t=t_{\mathrm{ret}}} , \quad (1)$$

where G is the gravitational constant, while M_j, $\mathbf{q}^{(j)}$ and γ_j, $j = 1, \ldots, N$, are the mass, the position vector and the Lorentz factor of the j–th source galaxy, the dot denoting derivative with respect to proper time along the worldline of the source.

As $h_{\mu\nu}$ depends on each source not only through its position, but also through its velocity, the latter has to be assigned in order that the model be defined. To this end we use Hubble's law as a phenomenological prescription. Making reference to a local chart with Lorentzian coordinates having as origin the center of mass of the considered localized system (the Coma cluster), for the velocities $\mathbf{q}_j \equiv \mathbf{q}^{(j)}$ of the external galaxies we thus require

$$\dot{\mathbf{q}}_j = \gamma_j^{-1} H_0 \mathbf{q}_j \ , \quad j = 1, \ldots, N \quad (2)$$

(the dot denoting now derivative with respect to the background Lorentzian time). Here, H_0 is the Hubble constant which, in our extremely simplified model, we take fixed at its present value.

---

[*] Electronic address: carati@mat.unimi.it

[†] Electronic address: sergio.cacciatori@uninsubria.it

[‡] Electronic address: galgani@mat.unimi.it

Notice that the Hubble assumption (2) has an essential impact on the size of the gravitational field of force. Indeed such a force contains a term (decreasing as $1/r^2$) proportional to the velocity of the source, and a term (decreasing as $1/r$) proportional to the acceleration of the source. Thus, estimating the acceleration through Hubble's law, the latter term actually doesn't depend on distance at all, so that the far matter is found to give the dominant contribution to the gravitational field of force. This situation is reminiscent of the way in which Mach's principle was dealt with in [3] (see page 102). The main difference being that in such a case, lacking Hubble's law, the velocities of the sources were neglected. Consequently, only the Newtonian, fast decaying, potential was considered, so that only the near matter, and not the far one, appeared to play a role.

So we address our attention to the dominating term, proportional to the acceleration of the source. Such a term, which we denote by $\mathbf{f}$, has the form

$$\mathbf{f} = \frac{4GH_0^2\,M}{c^2}\,\mathbf{u}\ , \qquad \mathbf{u}(N) = \sum_{j=1}^{N} \frac{\mathbf{q}_j}{|\mathbf{q}_j|}\ , \tag{3}$$

where the masses of the galaxies were all put equal to a common value M, and the Lorentz factors γ_j were put equal to 1, for the reasons to be illustrated later. Notice the extremely simple nature of this force per unit mass (or acceleration). Apart from a multiplicative factor, such a force is just the sum of all the unit vectors pointing to each of the external galaxies. Actually, our attention was addressed to the component of such a force $\mathbf{f}$ along a given direction. Such a component will be simply denoted by f, and the corresponding component of $\mathbf{u}$ by u.

Having determined the quantity of interest (f or u), we come now to our second main point, namely the problem of how to describe the distribution of the external galaxies. It is immediately seen that u exactly vanishes (at any point) if the external matter is described as a continuous medium with a spherically symmetric density. This should be expected, in agreement with Birkhoff's theorem. So we take a different point of view, analogous to the probabilistic one introduced by Chandrasekhar and von Neumann (see the review [4]) in the problem of estimating the vector sum of the Newtonian forces exerted on a star by the surrounding ones. In such an approach, the external sources are conceived as point particles, and the place of the matter density of the continuum case is now taken by a probability density for the position of a galaxy. From such a point of view, the previous result (the vanishing of u for a spherically symmetric matter density), now reads as the vanishing of the mean value of u for a spherically symmetric probability density of the position of a galaxy.

We thus come to an estimate of the variance of the force f (or of u). It will be seen that the result depends on the further assumptions one introduces concerning the spatial distribution of the external galaxies. Assume first that the positions $\mathbf{q}_j$ of the N galaxies are independent random variables, uniformly distributed with respect to the Lebesgue measure. Then the sum defining u is found to grow as $\sqrt{N}$. This indeed is just a consequence of the central limit theorem, because in such a case u is the sum of N independent identically distributed random variables which turn out to have zero mean and a finite variance. By the way, such a result is the analogue of that obtained by Chandrasekhar and von Neumann for the case of Newtonian forces, the only difference being that in their case the variance is infinite (due to the divergence of Newton's force at zero distance, a property which plays no role in our case). For what concerns the corresponding estimate of the virial, one easily sees that with the present assumption the estimate is by far too small to account for the observations, just because the considered sum behaves as $\sqrt{N}$ rather than as N (see later).



FIG. 1: Histogram of the random variable u, which is proportional to the force per unit mass due to N external galaxies. The histogram was computed with $10,000$ samples of $N = 512,000$ galaxies, by counting the fraction of times the value u belongs to a given interval of width $\sqrt{N}$.

So we modify the previous assumption, and consider the case in which the probability density is fractal [5] (see also [6]). This means first of all that the positions of the galaxies are no more independently distributed, so that u is no more constrained to grow as $\sqrt{N}$, and can instead have a faster growth, as required by the observations. However, the analytical computation of the probability distribution of the field of force now becomes a quite nontrivial task, with respect to the much simpler case considered by Chandrasekhar and von Neumann. So we are forced, at least provisionally, to investigate the problem by numerical methods.

We proceeded as follows. In order to estimate (at the origin of the coordinates) the sum defining u, the positions of the N galaxies are extracted (with the method



FIG. 2: The variance σ_u^2 of u versus the number N of galaxies in log–log scale. The dashed line is the curve $\sigma_u^2 = 0.2\ N^2$.

described in [5]) in such a way that the mass distribution has a fractal dimension, precisely the fractal dimension 2. The corresponding histogram, obtained through 10,000 samples of configurations of the galaxies, is shown in Fig. 1 for $N = 512,000$.

We then study the dependence of u on the number N of external galaxies, which was made to vary in the range $1000 \leq N \leq 512,000$, with the density kept constant. This means that the positions of the N points were taken to lie inside a cutoff sphere whose volume was made to increase as N. For the values of N investigated, the corresponding radius turns out to be so small with respect to the present horizon, that the Lorentz factors γ could altogether be put equal to 1 (as was previously assumed), and more in general the special relativistic character of our model was actually justified.

The mean of u turns out to practically vanish for all N, while its variance σ_u^2 is found to grow as N^2 (actually, as $0.2\,N^2$), rather than as N, as occurs in the uniform case. This is shown in Fig. 2. We thus conclude that the standard deviation σ_f of the component of the force per unit mass along a direction is proportional to N, being given by

$$\sigma_f \simeq \sqrt{0.2}\ \frac{4GH_0^2}{c^2}\ MN\ = \sqrt{0.2}\ \frac{4G}{R_0^2}\ MN\ . \tag{4}$$

(where $R_0 = c/H_0$ is the present horizon).

We now take such a result, which was obtained for extremely small values of N, and extrapolate it up to the present horizon $R_0 = c/H_0$, i,e., we insert in formula (4) the actual value of N, so that the quantity MN can be identified with the total visible mass of the Universe.

Concerning the total visible mass MN of the Universe, one can write

$$MN = \frac{4}{3}\pi\ \rho_{\rm eff}\ R_0^3\ , \tag{5}$$

with a suitable effective density $\rho_{\rm eff}$. It is rather easily shown (see later) that a quite natural consistency condition of our model leads to the estimate

$$\rho_{\rm eff} \simeq \frac{1}{4}\,\frac{3H_0^2}{8\pi G} \simeq 5\rho_0\ , \tag{6}$$

where $\rho_0 = \Omega_0\,\big(3H_0^2\big)/\big(8\pi G\big)$, with $\Omega_0 \simeq 0.05$, is the actual density. Inserting this in (4) one gets $\sigma_f \simeq 0.2\ cH_0$.

On the other hand, if a random variable f has zero mean and a finite variance σ_f^2, with great probability its modulus will take on values very near to its standard deviation σ_f. In such a sense we may say to have found

$$|f| \simeq 0.2\,cH_0\ , \tag{7}$$

which perhaps constitutes the main result of the present work. Namely, in our oversimplified model within the fractal hypothesis, the force per unit mass, i.e., the acceleration, exerted by the far matter on a test particle, is found to have a value of the order of cH_0, which is the one that is needed in most cases in which the presence of a dark matter is advocated. Notice that the assumption of a uniform, rather than fractal, distribution of matter would lead instead to $|f| \simeq cH_0/\sqrt{N}$, i.e., essentially to $f \simeq 0$. Namely, without the fractal hypothesis the Zwicky procedure of neglecting at all the gravitational contribution of the external matter, would be justified.

We can now apply our estimate to the case of the virial theorem for a cluster of galaxies. To this end, we have first of all to assume a property that might in principle be checked, namely, that the forces at sufficiently separated points are uncorrelated. It is then conceivable that locally, in some regions, such a random field of force may form patterns of a central type, attractive towards a center. This is obviously equivalent to admit that locally, in such special regions, the external far matter produces a pressure.

Let us recall that, according to the virial theorem, for a confined system of n particles one has

$$\overline{\sigma_v^2} = -\overline{\mathcal{V}}/n\ ; \tag{8}$$

here, $\sigma_v^2 = (1/n)\sum_i v_i^2$ is the variance of the velocity distribution of the galaxies of the cluster, whereas $\mathcal{V} = \sum_i \mathbf{f}_i\cdot\mathbf{x}_i$ is called the virial of the forces (per unit mass), $\mathbf{x}_i$ denoting the position vector of the i–th internal particle with respect to the center of mass of the cluster, and $\mathbf{f}_i$ the force per unit mass acting on it, while overline denotes time–average,

So we have to estimate the quantity $\sum_{i=1}^n \overline{\mathbf{f}_i\cdot\mathbf{x}_i}$. In the conditions we have assumed, all terms of such a sum can be taken to be equal, so that we just have to estimate one of them. We can take $\overline{\mathbf{f}_i\cdot\mathbf{x}_i} \simeq -|\bar{f}|\ \overline{|\mathbf{x}_i|}$, with $\overline{|\mathbf{x}_i|} \simeq L/4$ where L is the diameter of the cluster, whereas for estimating the modulus of the force in the direction of the center one can make use of (7). So for the velocity variance one gets

$$\overline{\sigma_v^2} \simeq 0.2\ \frac{cH_0L}{4}\ , \tag{9}$$

where L is the linear dimension of the cluster. In the case of Coma, one thus finds a value $\simeq 6 \cdot 10^5 \mathrm{km}^2/\mathrm{sec}^2$, which is very near to the value $5 \cdot 10^5 \mathrm{km}^2/\mathrm{sec}^2$ reported by Zwicky.

Notice the linear dependence on L in the formula (9). In this connection one may point out that, if the external force were smooth, by a Taylor expansion about the origin one would have $\mathbf{f}_i$ proportional to $\mathbf{x}_i$, and this would lead to a virial (and thus also a velocity variance) proportional to L^2 rather than to L. Instead, the observations seem to require a proportionality to L. Apparently, this was first pointed out by Kazanas and Mannheim [7], in a paper in which some data were reported in a range of L covering five orders of magnitude (see [7], Fig. 2, page 539). This property is also confirmed by a dimensional analysis. Indeed, with the parameters entering the problem, the square of a velocity can be formed only as c^2, or as cH_0L or as $(H_0L)^2$. But the first term is by far too large, the last term by far too small, while the term linear in L is indeed about of the correct order of magnitude.

We finally show how the estimate (6) for the effective density, namely, $\rho_{\mathrm{eff}} \simeq 5\rho_0$, is obtained. To this end one makes reference to the metric $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ with $h_{\mu\nu}$ defined by (1), and to the corresponding mean metric obtained by averaging with respect to the considered probability density for the positions of the galaxies. Denoting the mean by $\langle\, . \rangle$, the mean metric is then found to be given by

$$ds^2 = \langle\, g_{\mu\nu} \,\rangle\, dx^\mu dx^\nu = (1-\alpha-3\beta)\, c^2dt^2 - (1+\alpha+\beta) dl^2$$

where $dl^2 = dx^2 + dy^2 + dz^2$ and

$$\alpha = \frac{2G}{c^2}\, \langle \sum_j \frac{M_j}{|\mathbf{q}_j|} \rangle \ , \quad \beta \lesssim \frac{4GH_0^2}{3c^4}\, \langle \sum_j M_j |\mathbf{q}_j| \,\rangle \qquad (10)$$

(the approximation of small velocities was used in the expression for β). Introducing an effective density ρ_{eff} such that

$$\langle \sum \frac{M_j}{|\mathbf{q}_j|} \rangle \simeq 4\pi\rho_{\mathrm{eff}}\, \frac{{R_0}^2}{2} \ , \quad \langle \sum M_j |\mathbf{q}_j| \,\rangle \simeq 4\pi\rho_{\mathrm{eff}} \frac{{R_0}^4}{4} \ ,$$

one gets

$$\alpha \simeq 4\pi G \rho_{\mathrm{eff}} {R_0}^2/c^2 \ , \quad \beta\ < (2/3)\alpha \ . \qquad (11)$$

The consistency condition is now the requirement that the expansion rate calculated with the mean metric does actually coincide with the rate that was introduced into the definition of the model. This condition takes the form

$$\frac{1}{2}\frac{d}{dt} \log \frac{1+\alpha+\beta}{1-\alpha-3\beta} = H_0 \ . \qquad (12)$$

On the other hand from (11), using $\dot{R}_0 = c$, one gets

$$\dot{\alpha} \simeq \frac{8\pi G}{c^2}\, \rho_{\mathrm{eff}}\, R_0 c \ , \quad \dot{\beta} \simeq \frac{2}{3}\dot{\alpha} \ .$$

With these expressions for $\dot{\alpha}$ and $\dot{\beta}$, the consistency condition (12) then becomes an algebraic one, which gives for ρ_{eff} a value that we have rounded off to (6).

Acknowledgments

We thank George Contopoulos, Christos Efthymiopoulos. Francesco Sylos Labini and Rudolf Thun for useful discussions.

[1] F. Zwicky, Helv. Phys. Acta **5**, 110 (1933); Astrophys. J. **86**, 217 (9137).

[2] Ya.B. Zeldovich, I.D. Novikov, *Stars and relativity* (Dover, New York 1971).

[3] A. Einstein, *The meaning of relativity* (Princeton U.P., Princeton 1922).

[4] S. Chandrasekhar, Rev. Mod. Phys. **15**, 1 (1943).

[5] B. Mandelbrot, *The fractal geometry of nature* (Freeman, New York 1977).

[6] F. Sylos–Labini, M. Montuori, and L. Pietronero, Phys. Rep. **293**, 61 (1998).

[7] D. Kazanas, P.D. Mannhein, in *After the first three minutes*, edited by S. Holt, C.L. Bennett, B.V. Trimble, AIP Conference Proceedings Vol. 222 (American Institute of Physics, New York 1991).